SMAP Science and Hydros Heritage

Dara Entekhabi


SMAP Mission Workshop Arlington, VA July 9-10, 2007



| Decadal Survey Panel | Cited Soil Moisture Applications |
|--|---|
| Water Resources and Hydrological Cycle | Floods and Drought Forecasts Available Water Resources Assessment Link Terrestrial Water, Energy and Carbon Cycles |
| Climate / Weather | Longer-Term and More Reliable Atmospheric Forecasts |
| Human Health and Security | Heat Stress and Drought Vector-Borne and Water-Borne Infectious Disease |
| Land-Use, Ecosystems, and Biodiversity | Ecosystem Response (Variability and Change) Agricultural and Ecosystem Productivity Wild-Fires Mineral Dust Production |

"...the SMAP mission is ready for "fast-track" towards launch as early as 2012, when there are few scheduled Earth missions. The readiness of the SMAP mission also enables gap-filling observations to meet key NPOESS community needs (soil moisture is "Key Parameter," see 4.1.6.1.6 in IORD-II Document)."

Page 4-43



Panel Recommendations:

TABLE 11.1 Candidate Missions in the Order Ranked

Water Resources and Hydrological Cycle Panel:

| Brief Description of Mission | Variables | Type of Sensor(s) | Coverage | Spatial Resolution | Frequency | Synergies with other panels | Related Planned or Integrated Missions (if any) |
|---|--|--|---------------------------------|-----------------------------------|--------------------|--|---|
| Soil Moisture and Freeze/ Thaw State | Surface freeze/thaw state; Soil moisture | L-band radar, radiometer | Global | 10 km (processed to 1-3 km) | 2-3 day revisit | Climate Weather | SMAP Aquarius |
| Surface Water and Ocean Topography | River and lake elevation; Ocean circulation | Radar altimeter, Nadir SAR interferometer, microwave radiometer, GPS receiver | Global (to ~82° latitude) | Several cm (vertical) | 3-6 days | Climate Ecosystems Health Weather | SWOT SMAP GPM NPP/ NPOESS |
| Snow and Cold Processes | Snow water equivalent; Snow depth; Snow wetness | SAR, Passive microwave radiometry | Global | 100 m | 3-15 days | Climate Ecosystems Weather | SCLP |


Land-Use, Ecosystems, and Biodiversity Panel:


Soil moisture is a key measurement for several disciplines, especially hydrology, where the primary discussion of this mission will be found. We strongly endorse a soil moisture experiment. In order to maximize the value of a soil moisture measurement for ecosystem science, it needs to resolve the time and space scales of variability relevant to ecosystem science. A temporal resolution (repeat sampling interval) of 3-5 days is needed to allow successful assimilation and inference of available water. This time interval is also critical for monitoring the development of plant water limitation and wet intervals associated with rapid and important soil activity. The spatial resolution required must correspond to scales of variability in terrestrial ecosystems, and in the soil moisture anomalies that affect them. This implies spatial resolution on the order of square kilometers to tens of square kilometers.

(Page 7-22)

Decadal Survey: "...delivery of flash-flood guidance to weather forecast offices are centrally dependent on the availability of soil moisture estimates and observations."

"SMAP will provide realistic and reliable soil moisture observations that will potentially open a new era in drought monitoring and decision-support."





Current: Empirical Soil Moisture Indices Using Rainfall and Air Temperature (Counties ~30 km)

SMAP Soil Moisture Observations at 10 km Resolution




Soil Moisture and Its Impact on Atmosphere


May 10: Clear with scattered cirrus

PBL Winds 5-7 m/s

May 18: 90 mm Rain

May 20: Clear sky & mild winds







CASES'97, BAMS (81), 2000.



Soil Moisture and Weather Forecasts



Chen et al., JAS (58) 2001.



"The strong motivation for this land data assimilation and land-monitoring space mission such as HYDROS is that the land states of soil moisture, soil ice, snowpack, and vegetation exert a strong control on ...the heating and moistening of the lower atmosphere...forecast of tomorrow's heat index, precipitation, and severe thunderstorm likelihood."

Louis Uccellini, NCEP

"The experience of the last ten years at ECMWF has shown the importance of soil moisture...Soil moisture is a major player on the quality of weather parameters such as precipitation, screen-level temperature and humidity and low-level clouds."

Anthony Hollingworth, ECMWF




Soil Moisture and Climate Prediction

Regions Where Soil Moisture Has Significant* Impact on Seasonal Precipitation Predictability Through Land-Atmosphere Coupling (Dots)

Product Retrieval Error Map (Blue Shades in % Volumetric)

Koster et al. 2003, J. Hydromet. 4[2])





*At Least 1 in 5 Summers Affected in Model Simulations




Response From Multiple Climate Models


Summer 1993 Rainfall Minus Summer 1988 Rainfall












Koster et al. Science, 305, 2004.





Global Change and Water Cycle/Resources

"...models suited for estimating first-order changes in atmospheric temperature profiles...[are] notoriously challenged with regards to reproducing and predicting changes in atmospheric wet processes..."

GEWEX News, 11(1)









Li et al., 2007: JGR (112)



Decadal Survey:

"...major source of uncertainty in assessing the carbon budget of the Earth system (... "missing carbon sink").... In boreal latitudes, the switching on and off of the land-atmosphere carbon exchange is coincident with the freeze/thaw transitions."






Frolking et al., Global Change Biol. 2, 1996.




Freeze/Thaw Science Requirements

State



JERS-1 L-band SAR landscape freeze-thaw classification




To resolve Carbon source/sink to within 0.05 [tons C ha⁻¹]*

2 km resolution 1-2 day refresh

0.7 dB accuracy



* 5 [g C m-2] over 100 days



Science Return: Global Views of Terrestrial Water Cycle State Variable: Soil Moisture Content and Its Freeze/Thaw State

Soil Moisture is a Variable That <u>Links</u> the Global Water, Energy, and Carbon Cycles

Applications Return: A New Era for the Capability to Predict Costly Natural Hazards (Extreme Weather, Floods, Droughts)

Initialization of the Soil Moisture State in Numerical Models Extends the Predictability of Processes Influenced by Surface Fluxes

National Defense Return: Global All-Weather Mapping Supports Battlespace Decision-Making and Force-Enhancement

Army Terrain Trafficability
Air Force Low-Level Fog, Dust and NWP
Navy Sea-Ice Edge and Age Assessment

Track-Record of User-Community Advocacy



| Tri-Service Flag Officers (DoD) | Military Applications |
|---|---|
| Army Research Laboratory (ARL) | Army Trafficability Application |
| National Centers for Environmental Prediction (NCEP) | Operational Weather Prediction |
| European Centre for Medium-Range Weather Forecasts | Operational Weather Prediction |
| Meteorological Service of Canada (MSC) | Operational Weather Prediction |
| NASA Seasonal to Interannual Prediction Project (NSIPP) | Seasonal Climate Prediction Application |
| NOAA Climate Prediction Center (NOAA CPC) | Seasonal Climate Prediction Application |
| GEWEX Americas Prediction Project (GAPP) | Regional Water Cycle Applications |
| National Drought Mitigation Center (NDMC) | Drought Hazard Mitigation Application |
| SAHRA - Sustainability of semi-Arid Hydrology and Riparian | Water Resources Application |
| Office of Global Carbon Studies | Carbon Cycle Science Application |
| U.S. Geological Survey (USGS) | Water Resources and Flood Application |
| Dartmouth Flood Observatory (DFO) | Flood Monitoring Application |
| Center for Hydrology, Soil Climatology, and Remote Sensing | Field Site and Outreach |
| Boston Museum of Science | Education and Public Outreach |
| Global Learning and Observations to Benefit the Environment | Education |
| | |



SMAP Primary Science Objectives

- Collect space-based measurements needed to retrieve estimates of soil moisture and freeze/thaw state, for at least 3 years, over global land regions where these factors are the primary environmental controls on surface-atmosphere fluxes of water and carbon.
- 2. Provide estimates of soil moisture in the top 5 cm of the surface with a relative error of 4-5% volumetric, at 3-day intervals, at 10-40 km spatial resolution (hydrometeorology to hydroclimate spatial scales)
- 3. Provide estimates of surface freeze/thaw transitions in Boreal zones with 2-3 day relative error, at 1-3 km spatial resolution (sufficient to resolve dominant landscape heterogeneity)
- 4. Record, calibrate, validate, publish, and archive science data records and calibrated geophysical data products for use by the scientific community
- 5. Validate a space-based measurement approach and analysis concept that can be used for future systematic soil moisture and freeze/thaw monitoring missions



Requirements Traceability

| Science Objectives | Scientific Measurement Requirements | Instrument Functional Requirements | Mission Functional Requirements |
|--|--|---|---|
| Enhance understanding of processes that link the water, energy and carbon cycles through monitoring the land hydrosphere state. Extend the capability of climate prediction models by providing land surface boundary data products with required resolution, global coverage, and revisit times. | Soil Moisture: ~±4% volumetric accuracy in top 2-5 cm for vegetation water content < 5 kg m ⁻² ; Hydrometeorology at ~10 km; Hydroclimatology at ~40 km | L-Band Radiometer: Polarization: V, H, U; Resolution: 40 km; Relative accuracy*: 1.5 K L-Band Radar: Polarization: VV, HH, HV; Resolution: 10 km; Relative accuracy*: 0.5 dB for VV and HH Constant incidence angle** between 35° and 50° | DAAC data archiving and distribution. Field validation program. Integration of data products into multisource land data |
| | Freeze/Thaw State: Capture freeze/thaw state transitions in integrated vegetation-soil continuum with two-day precision, at the spatial scale of land-scape variability (~3 km). | L-Band Radar: Polarization: HH; Resolution: 3 km; Relative accuracy*: 0.7 dB (1 dB per channel if 2 channels are used); Constant incidence angle** between 35° and 50° | assimilation. |
| | Sample diurnal cycle at consistent time of day (6am/6pm); Global, ~3 day revisit; Boreal, ~2 day revisit | Swath Width: ~1000 km Minimize Faraday rotation (degradation factor at L-band) | Orbit: 670 km, circular, polar, sun-synchronous, ~6am/pm equator crossing |
| | Observation over minimum of three annual cycles | Minimum three-year mission life | Three year baseline mission*** |

^{*} Includes precision and calibration stability

^{***} From end of 30 days post-launch observatory check-out



^{**} Defined without regard to local topographic variation

Summary

- SMAP Mission Data Will be Utilized Across Broad Range of Earth Science and Application Areas.
- 2. High-Resolution and High-Revisit Data Will be Transformational for Several Earth System (Water, Energy and Carbon Cycles) and Operational Applications.
- 3. Advocacy and Partnership History Stand on Strong Foundations.
- 4. Mission Goals and Requirements-Traceability Are Advanced.
- 5. Hydros Science Heritage is Substantial and Well-Documented.
 - a. Level-1 Requirements and Mission Success Criteria Document
 - b. Level-1 Science Requirements Document
 - c. Ten Data Products Algorithm Roadmaps (pre-ATBD)